THEF-NIJM 94.04

Partial Wave Analyses of the *pp* data alone and of the *np* data alone*

R.A.M.M. Klomp, J.-L. de Kok, M.C.M. Rentmeester, Th.A. Rijken, and J.J. de Swart
Institute for Theoretical Physics, University of Nijmegen, Nijmegen, The Netherlands

Abstract

We present results of the Nijmegen partial-wave analyses of all NN scattering data below $T_{\rm lab} = 500$ MeV. We have been able to extract for the first time the important np phase shifts for both $I = 0$ and $I = 1$ from the np scattering data alone. This allows us to study the charge independence breaking between the pp and np $I = 1$ phases. In our analyses we obtain for the pp data $\chi^2_{\rm min}/N_{\rm df} = 1.13$ and for the np data $\chi^2_{\rm min}/N_{\rm df} = 1.12$.

Typeset using REVTEX

*Contribution to the XIVth International Conference on Few-Body Problems in Physics, May 26–31, 1994, Williamsburg, VA. Published in the proceedings: AIP Conf. Proc. **334**, 367–369 (1995)

I. INTRODUCTION

The last 15 years the Nijmegen group has been working on partial-wave analyses (PWA) of the NN-scattering data for energies below $T_{\rm lab} = 350$ MeV. In these analyses, the pp data furnish the $I = 1$ phase shifts [1] in the lower partial waves with $J \leq 4$. It has been customary to do these low energy PWA up to $T_{\rm lab} = 350$ MeV, despite of the fact that the inelasticities set in at $T_{\rm lab} \simeq 280$ MeV. It can be shown, however, that up to $T_{\rm lab} = 350$ MeV the inclusion of inelasticities in pp scattering improves the total $\chi^2 \simeq 1787$ with only about 1, so neglecting the inelasticities is totally warranted.

In the PWA of all the np data below $T_{\rm lab} = 350$ MeV, it has always been impossible to determine all important phase shifts, when only the np data were used. The standard procedure has therefore been to take the $I = 1$ phases (except the ${}^1S_0$) over from the pp analyses, with or without corrections for Coulomb, and np and $\pi^0\pi^\pm$ mass difference effects. The ${}^1S_0$ np phase shift was always searched for. It was found that there is a definite charge independence breaking in these ${}^1S_0$ phases. An attempt by us, to extract all the lower $I = 0$ and $I = 1$ phase shifts in an analysis of all np data below $T_{\rm lab} = 350$ MeV failed. However, it was possible to determine the np ${}^3P$ waves, when the higher partial waves were taken over from the pp analysis.

Recently, the Nijmegen group extended the pp PWA to energies so far above the pion production thresholds, that the inclusion of inelasticities was necessary. A preliminary version of such a PWA for all pp data with energies below $T_{\rm lab} = 500$ MeV has already been presented [2].

When the np PWA was extended to energies up to $T_{\rm lab} = 500$ MeV, it turned out to be possible to determine uniquely all the important partial waves. Comparing then with the analogous pp analysis gives indications for possible charge independence breaking effects in other waves besides the ${}^1S_0$ waves.

II. THE METHOD OF ANALYSIS

In the np as well as in the pp partial wave analyses, the various phases are obtained by solving the relativistic Schrödinger equation with a well-known long-range potential $V_{\rm L} = V_{\rm NUC} + V_{\rm EM}$ for $r \geq b = 1.4$ fm. This $V_{\rm L}$ contains the electromagnetic interaction (such as Coulomb, magnetic moment, and vacuum polarization), the OPEP tail, and the rest of the long range NN interaction as given by the Nijmegen potential [3].

For $r < b$ the short-range interaction is described by an energy-dependent, square-well potential

$$V_{\rm S} = V_{\rm R} - iV_{\rm I} \ .$$

For $V_{\rm R}$ we follow the same procedure as in the Nijmegen PWA and write

$$V_{\rm R} = \sum_{n=0}^{N} a_n (k^2)^n \ .$$

For the imaginary part of the potential we take

$$V_{\rm I} = (k^2 - k^2_{\rm thr})^n V \cdot \theta(E - E_{\rm thr}) \ .$$